


PRACTICE A

Key

4. Identify any holes, vertical asymptotes, horizontal asymptotes, and slant asymptotes of each graph. If they do not exist write **none** in the space provided. Accurately graph each rational function.

a. $f(x) = \frac{x}{x-1}$

Domain $x \neq 1$ $(-\infty, 1) \cup (1, \infty)$ Hole(s) none



Vertical Asymptotes $x = 1$

X-intercepts $(0, 0)$ Horizontal Asymptotes $y = 1$

Set $y = 0$

$0 = x$

BOSCO

Y-intercepts $(0, 0)$ Slant Asymptote none

$x = 0$

7. If you have a BOTNO, then there is no HA. Instead, there is a slant or oblique asymptote.

TO FIND THE EQUATION OF A SLANT ASYMPTOTE, you simply perform the division that is left in the working equation. Any remainder is NOT part of the slant asymptote.

$$f(x) = \frac{x^2 + 2x - 5}{x - 3}$$

Example: BOTNO

SLANT

$$(x^2 + 2x - 5) \div (x - 3)$$

$$\begin{array}{r} 3 \overline{) 1 2 - 5} \\ \underline{3 15} \\ 1 5 10 \end{array}$$


STEPS:

1. DOMAIN RESTRICTIONS $x \neq 3$
2. HOLES?
3. x intercept
4. y INTERCEPT
5. VA
6. HA

$$y = 1x + 5 = \text{slant asy.}$$

Domain $(-\infty, 3) \cup (3, \infty)$

The graph can only cross x and y axes at the intercepts you've found.
VA's are fences that the graph cannot cross.
Graphs hug asymptotes



$$x = 4 \\ f(x) = \frac{16 + 8 - 5}{1} = \frac{19}{1}$$

$$V.A. x = 3$$

$$\text{Slant } y = x + 5$$

$$(0, 5/3)$$

$$(-1, 0)$$



$$0 = x^2 + 2x - 5$$

$$x = \frac{-b \pm \sqrt{b^2 + 4ac}}{2a}$$

$$x = \frac{-2 \pm \sqrt{4 + 20}}{2}$$

$$x = \frac{-2 \pm \sqrt{24}}{2} = \frac{-2 \pm 2\sqrt{6}}{2}$$

$$x = -1 \pm \sqrt{6}$$

b. $f(x) = \frac{x^2 - x - 12}{x - 4}$

$y = \frac{(x-4)(x+3)}{(x-4)}$


Key

Domain $x \neq 4$ Hole(s) $(4, 7)$

$(-\infty, 4) \cup (4, \infty)$

Vertical Asymptotes none

line with hole



X-intercepts $(-3, 0)$ Horizontal Asymptote —

$y = x + 3$


Y-intercepts $(0, 3)$ Slant Asymptote —

c. $f(x) = \frac{x-3}{x^2 - 5x + 6} = \frac{(x-3)}{(x-3)(x-2)} = \frac{1}{x-2}$

Domain $x \neq 3, 2$ Hole(s) $(3, 1)$

$(-\infty, 2) \cup (2, 3) \cup (3, \infty)$

Symmetry — Vertical Asymptotes $x = 2$



X-intercepts none Horizontal Asymptotes $y = 0$

BOBO

Y-intercepts $0, -1/2$ Slant Asymptotes —


d. $f(x) = \frac{(x+2)(x-3)}{x(x-4)^2(x+2)}$

$\frac{-5}{-2(36)} = \frac{5}{72}$

Domain $x \neq 0, 4, -2$ Hole(s) $(-2, 5/72)$


$(-\infty, -2) \cup (-2, 0) \cup (0, 4) \cup (4, \infty)$

Vertical Asymptotes $x = 0$ $x = 4$



X-intercepts $3, 0$ Horizontal Asymptotes $y = 0$

BOBO




Y-intercepts none Slant Asymptotes none
 $-3/$

e. $f(x) = \frac{x}{x^2 - 4}$

Domain $x \neq 2, -2$ Hole(s) —


$(-\infty, -2) \cup (-2, 2) \cup (2, \infty)$

Symmetry — Vertical Asymptotes $x = 2$ $x = -2$



X-intercepts $(0, 0)$ Horizontal Asymptotes $y = 0$

BOBO



Y-intercepts $(0, 0)$ Slant Asymptotes —

Suggested

Homework: p. 342 #'s 2-8 (even), 9-14 (all), 38-42 (even), 52-76 (every other even)